



Individualized Fortification of Breast Milk for Preterm Infants

Miami Neonatology 2018—42nd Annual International Conference

Supplemental Questions

Successful completion of this activity is achieved by individually reflecting on, or discussing as a group, the following questions and their implications in your practice.

1. What are some challenges in your practice for patients using breast milk (mother or donor milk) in preterm infants?

See bottom of page 3, top of page 4 of the transcript.

2. How do you determine whether adequate enteral nutrients are reaching your preterm infant patients?

See bottom of page 6 of the transcript.

3. How do you select the right nutritional strategy for your patients?

See page 11, below slide 22 of the transcript.

4. In your opinion, how have fortified infant formulas progressed over the last decade and what challenges remain?

See page 5 of the transcript.

5. What characteristics should the optimum composition for an infant feeding fortifier have to meet the needs of your preterm infant patients?

See bottom of page 11, below slide 22 of the transcript.

Suggested Reading and References

Agostoni C, Buonocore G, Carnielli VP, et al. Enteral nutrient supply for preterm infants: commentary from the European Society of Paediatric Gastroenterology, Hepatology and Nutrition Committee on Nutrition. *J Pediatr Gastroenterol Nutr.* 2010;50(1):85-91. doi:10.1097/MPG.0b013e3181adaee0

Arslanoglu S, Moro GE, Ziegler EE. Adjustable fortification of human milk fed to preterm infants: does it make a difference? *J Perinatol.* 2006 Oct;26(10):614-621.

Cooke RJ, Ainsworth SB, Fenton AC. Postnatal growth retardation: a universal problem in preterm infants. *Arch Dis Child Fetal Neonatal Ed.* 2004;89(5):F428-30.

Individualized Fortification of Breast Milk for Preterm Infants

Cooke RJ, Embleton ND. Feeding issues in preterm infants. *Arch Dis Child Fetal Neonatal Ed.* 2000 Nov;83(3):F215-8.

Corvaglia L, Aceti A, Paoletti V, et al. Standard fortification of preterm human milk fails to meet recommended protein intake: Bedside evaluation by Near-Infrared-Reflectance-Analysis. *Early Hum Dev.* 2010;86(4):237-40. doi:10.1016/j.earlhumdev.2010.04.001

Ehrenkranz RA, Younes N, Lemons JA, et al. Longitudinal growth of hospitalized very low birth weight infants. *Pediatrics.* 1999;104(2 Pt 1):280-289.

Embleton NE, Pang N, Cooke RJ. Postnatal malnutrition and growth retardation: an inevitable consequence of current recommendations in preterm infants? *Pediatrics.* 2001;107(2):270-273.

Embleton ND. Optimal protein and energy intakes in preterm infants. *Early Hum Dev.* 2007;83(12):831-837.

Fusch G, Mitra S, Rochow N, Fusch C. Target fortification of breast milk: levels of fat, protein or lactose are not related. *Acta Paediatr.* 2015;104(1):38-42. doi:10.1111/apa.12804

Gebauer CM, Ackermann B, Ristau J, et al. Standard Fortification (STD) Versus Target Fortification (TFO) in Very low Birthweight Infants: Effect On Growth and Amino Acid Profile; The LEIPZIG Trial (e-Poster EAPS8-1096).

Henriksen C, Westerberg AC, Rønnestad A, et al. Growth and nutrient intake among very low birth weight infants fed fortified human milk during hospitalization. *Br J Nutr.* 2009, 102, 1179–1186. doi:10.1017/S0007114509371755

Kashyap S, Ohira-Kist K, Abildskov K, et al. Effects of quality of energy intake on growth and metabolic response of enterally fed low-birth-weight infants. *Pediatr Res.* 2001;50(3):390-397.

Kashyap S, Towers HM, Sahni R, Ohira-Kist K, Abildskov K, Schulze KF. Effects of quality of energy on substrate oxidation in enterally fed, low-birth-weight infants. *Am J Clin Nutr.* 2001;74(3):374-380.

Lau CSM, Chamberlain RS, Sun S. Less invasive surfactant administration reduces the need for mechanical ventilation in preterm infants: A meta-analysis. *Glob Pediatr Health.* 2017; 4: 2333794X17696683. doi:10.1177/2333794X17696683

Madore LS, Bora S, Erdei C, Jumani T, Dengos AR, Sen S. Effects of donor breastmilk feeding on growth and early neurodevelopmental outcomes in preterm infants: An observational study. *Clin Ther.* 2017;39(6):1210-1220. doi:10.1016/j.clinthera.2017.05.341

McLeod G, Sherriff J, Hartmann PE, Nathan E, Geddes D, Simmer K. Comparing different methods of human breast milk fortification using measured v. assumed macronutrient composition to target

Individualized Fortification of Breast Milk for Preterm Infants

reference growth: a randomised controlled trial. *Br J Nutr.* 2016;115(3):431-439.

doi:10.1017/S0007114515004614

Micheli JL, Schutz T. Protein. In: Tsang RC, Lucas A, Uauy R, Zlotkin S. eds. *Nutritional needs of the Preterm Infant: Scientific Basis and Practical Guidelines.* Baltimore, MD: Williams & Wilkins, 1993: 29–46. 44.

Montjoux-Régis N, Cristini C, Arnaud C, Glorieux I, Vanpee M, Casper C. Improved growth of preterm infants receiving mother's own raw milk compared with pasteurized donor milk. *Acta Paediatr.* 2011;100(12):1548-1554. doi:10.1111/j.1651-2227.2011.02389.x

O'Connor DL, Gibbins S, Kiss A, et al. Effect of supplemental donor human milk compared with preterm formula on neurodevelopment of very low-birth-weight infants at 18 months: A randomized clinical trial. *JAMA.* 2016 Nov 8;316(18):1897-1905. doi:10.1001/jama.2016.16144

Polberger S. New approaches to optimizing early diets. *Nestle Nutr Workshop Ser Pediatr Program.* 2009;63:195-204; discussion 204-208, 259-268. doi:10.1159/000209982

Quigley M, Embleton ND, McGuire W. Formula versus donor breast milk for feeding preterm or low birth weight infants. *Cochrane Database Syst Rev.* 2018;6:CD002971. doi:10.1002/14651858.CD002971.pub4

Rochow N, Fusch G, Choi A, et al. Target fortification of breast milk with fat, protein, and carbohydrates for preterm infants. *J Pediatr.* 2013;163(4):1001-1007. doi:10.1016/j.jpeds.2013.04.052

Rozance PJ, Hay WW. Describing hypoglycemia—definition or operational threshold? *Early Hum Dev.* 2010; 86(5): 275–280. doi:10.1016/j.earlhumdev.2010.05.002

Sauer CW, Boutin MA, Kim JH. Wide Variability in caloric density of expressed human milk can lead to major underestimation or overestimation of nutrient content. *J Hum Lact.* 2017;33(2):341-350. doi:10.1177/0890334416672200

Simmer K. Human milk fortification. *Nestle Nutr Inst Workshop Ser.* 2015;81:111-121. doi:10.1159/000365808

Stephens BE, Walden RV, Gargus RA, et al. First-week protein and energy intakes are associated with 18-month developmental outcomes in extremely low birth weight infants. *Pediatrics.* 2009;123(5):1337-1343. doi:10.1542/peds.2008-0211

Weber A, Loui A, Jochum F, Bühner C, Obladen M. Breast milk from mothers of very low birthweight infants: variability in fat and protein content. *Acta Paediatr.* 2001;90(7):772-775.